ZAMP Bionews

« Journal Of Clinical Investigation Table Of Contents: Nov. 20, 2008

UC San Diego Announces Collaboration With Johnson & Johnson Pharmaceutical

Research & Development »

Output

Description

Research & Development »

Research & Development »

Research & Development »

Output

Description

Description

Research & Development »

Output

Description

Research & Development »

Description

Descrip

Visual Illusions: Barrow Scientists Solve 200-Year-Old Scientific Debate

Biotechnology Companies

Access Company Profiles, Find Sales Leads & Build Lists. Free Trial! www.selectory.com

Clinical Cancer Trials

Learn more about innovative cancer treatment options provided by CTCA.

CancerCenter.com

New Clinical Trial

Clinical trial for patients with glioblastoma (GBM) brain tumors www.novocuretrial.com

Ads by Google

Neuroscientists at Barrow Neurological Institute at St. Joseph's Hospital and Medical Center have discovered a direct link between eye motions and the perception of illusory motion that solves a 200-year-old debate.

Stephen Macknik, PhD, director of the Laboratory of Behavioral Neurophysiology; Susana Martinez-Conde, PhD, director of the Laboratory of Visual Neuroscience; Xoana G. Troncoso, PhD; and Jorge Otero-Millan; conducted a study

based on the Enigma painting, a visual illusion in which rotational motion is seen within a stationary image. The artwork has been at the center of a debate over whether the brain or the eye is behind the perception of illusory motion.

Dr. Martinez-Conde's laboratory recently discovered that microsaccades, a small, unconscious eye movement that occurs when humans fixate their eyes, are critical to normal vision. The team of scientists conducted the Enigma study to see if microsaccades are also behind the perception of this illusion. Based on their study, the hypothesis suggesting the illusion originates solely in the brain was ruled out.

Participants in the study observed the Enigma illusion while their eye movements were simultaneously recorded with high precision cameras. Microsaccade rates increased before the illusionary motion sped up and decreased before the motion slowed, revealing a direct link between the eye movements and the illusion.

"We have discovered that this illusion originates with eye movements and not solely the brain as previously thought," says Dr. Martinez-Conde. "The findings from the study could help design future prosthetics for patients with brain damage or brain lesions that affect the perception of motion."

Source:: Carmelle Malkovich

St. Joseph's Hospital and Medical Center

Share / Save ⇒

Tags: barrow neurological institute, brain damage, brain lesions, eye movement, eye movements, malkovich, motions, neurophysiology, neuroscientists, normal vision, phd director, precision cameras, prosthetics, s hospital, st joseph, stationary image, troncoso, visual illusion, visual neuroscience

Related posts

- Trigger For Brain Plasticity Identified: Signal Comes, Surprisingly, From Outside The Brain
- Trigger For Brain Plasticity Identified Signal Comes, Surprisingly, From Outside The Brain
- New Approach To Detect Early Progression Of Brain Tumors Identified
- Imaging Study Reveals Battle Between Lure Of Reward And Fear Of Failure
- Full Recovery From Head Injury Unlikely